

REFERENCES

- ANONYMOUS, (1915) : XXXII—Decades Kewensis Plantarum Novarum in Herbario Hortii Regnii Conservatarum Decas LXXXVI Kew, Bull. 1915: 344-350.
- BACKER, C. A. & BAKHUIZEN VAN DEN BRINK, R. C. (1965) : Flora of Java. vol. 2. Netherlands.
- BEDDOME, R. H. (1871) : Flora Sylvatica of Southern India. vol. 1. Madras.
- (1874) : Icones Plantarum Indiae Orientalis Madras
- CHOUDHURI, A. B. (1967) : Studies on the Meliaceae of India and neighbouring countries. Bull. Bot. Soc. Bengal 21 (1) 1: 1-24.
- GAMBLE, J. S. (1915) : Flora of the Presidency of Madras. vol. 1. London.
- (1918) : Notes on the Flora of Madras. Kew Bull. 1918: 222-228.
- HARMS (1940) : Family Meliaceae in Engler & Prantl. Pflanzenfamilien (ed. 2) 19b: 138-148, Berlin.
- WIGHT, R. & ARNOTT, W. (1834) : Prodromus Flora Peninsulae Indiae Orientalis. London.

39. TWO NEW COMBINATIONS OF THE GENUS *ASEMANTHIA* (STAPF) RIDLEY (RUBIACEAE)

Mussaenda Linn., one of the established genera of Rubiaceae with a number of species, is well characterised by one of the calyx lobes being usually deciduous and forming a large petaloid white or coloured leaf like structure. A closely related genus *Acranthera* Arn. ex Meissn. is referred by the more herbaceous form of the plants than the typical *Mussaenda* Linn. but was merged to *Mussaenda* Linn. by Baillon (1880) while Schumann (1891) maintained it as a distinct species. Later Stapf (1894) recorded it as a well marked genus after the exclusion of the species like *Acranthera maingayi* Hk. f. and *A. griffithii* Hk. f. Simultaneously he (l.c.) has proposed a new Section *Asemanthia* Stapf under *Mussaenda* Linn. considering the taxa of *Acranthera maingayi* Hk. f., *A. griffithii* Hk. f. and *Mussaenda mutabilis* Hemsl. which differ in the bushy straggly growth of the plants, absence of enlarged calyx-lobes and finally the long corolla tube. Ridley (1940) has elevated it to a distinct genus with distribution to Malay Peninsula and the islands of Borneo but nearer to Indian *Mussaenda* Linn.

His treatment is as under:

Asemanthia maingayi (Hk. f.) Ridley in Kew Bull. 1940:600.1940; Basionym: *Acranthera maingayi* Hk. f. in Fl. Brit. Ind. 3:92. 1880; Synonym: *Mussaenda mutabilis* Hemsl. in Hook. f. Ic. Pl. 18: t. 1718, 1887; *Mussaenda maingayi* (Hk. f.) Stapf in Trans. Linn. Soc. (Sr. 2) 4: 172. 1894.

King (1903) has described a new variety *hirsuta* under *Mussaenda mutabilis* Hemsl. on the basis of densely velvety nature of stem and under surface of leaves especially along the nerves. Ridley (1923) has also described one more variety *montana* under this same taxon based on dwarf compact nature of the plant with smaller leaves. On examination of these varieties we suggest as *Mussaenda mutabilis* Hemsl. belongs to the genus *Asemanthia* (Stapf) Ridley, a new combinations for the two taxa as:

Asemanthia maingayi (Hk. f.) Ridley var. *hirsuta* (King) Sinha et Mitra comb. nov.; Basionym: *Mussaenda mutabilis* Hemsl. var. *hirsuta* King in Jour. Asiat. Soc. Beng. 72(2): 182. 1903.

Syntype: Perak, King's collector 225 (CAL); Malacca, Derry 253 (CAL).

Asemanthia maingayi (Hk. f.) Ridley var.
montana (Ridley) Sinha et Mitra comb. nov.:

Basionym: *Mussaenda mutabilis* Hemsl. var.
montana Ridley in Fl. Mal. Peninsula 2:58.
1923.

Type: Malacca, Mount Ophir, Padang Batu,
Ridley 3215 (CAL).

BOTANICAL SURVEY OF INDIA,
INDIAN BOTANIC GARDEN,
HOWRAH-711 103,
February 22, 1980.

ACKNOWLEDGEMENTS

We thank to Mr. K. Rammurthy, Dr. G. G. Maity and Mr. G. S. Giri for valuable suggestions and help.

A. K. SINHA
B. MITRA

REFERENCES

- BAILLON, H. E. (1880): Rubiaceae in *Histoire des Plantes* 7: 257-546. Paris.
- KING, G. (1903): Materials for a flora of Malayan Peninsula XIV (*Mussaenda*). *Journ. Asiat. Soc. Beng.* 72(2): 181-187
- RIDLEY, H. N. (1923): The Flora of the Malaya Peninsula 2: 58-59. L. Reeve & Co., London.
- (1940): Notes on some Malayan Rubiaceae. *Kew Bull.* 1940: 593-613.
- SCHUMANN, K. (1891): Rubiaceae in A. Engler and K. Prantl's *Die Natürlichen Pflanzenfam.* 4 (4): 1-156. Leipzig.
- STAPF, O. (1894): On the flora of Mount Kinabalu in North Borneo. *Trans. Linn. Soc.* (Sr. 2) 4: 69-263.